

TABLE OF CONTENTS

<u>Section</u>	<u>Page</u>
I. Environmental Consequences of Long-Term Repository Performance	I-1
I.1 Introduction	I-1
I.2 Total System Performance Assessment Methods and Models	I-2
I.2.1 Features, Events, and Processes	I-4
I.2.1.1 Tectonic Activity	I-6
I.2.1.2 Erosion/Denudation	I-6
I.2.1.3 Periglacial Effects	I-7
I.2.1.4 Glacial and Ice Sheet Effects	I-8
I.2.1.5 Hydrostatic Pressure on Container	I-8
I.2.1.6 Soil and Sediment Transport	I-8
I.2.1.7 Seismic Damage to Waste Packages	I-9
I.2.2 Unsaturated Zone Flow	I-9
I.2.3 Engineered Barrier System Environments	I-11
I.2.4 Waste Package and Drip Shield Degradation	I-14
I.2.5 Waste Form Degradation	I-15
I.2.6 Engineered Barrier Transport	I-16
I.2.7 Unsaturated Zone Transport	I-16
I.2.8 Saturated Zone Flow and Transport	I-17
I.2.8.1 Saturated Zone Flow	I-18
I.2.8.2 Saturated Zone Transport	I-18
I.2.9 Biosphere	I-19
I.2.10 Volcanism	I-20
I.2.11 Human Intrusion	I-21
I.2.12 Nuclear Criticality	I-21
I.2.13 Atmospheric Radiological Consequences	I-21
I.3 Inventory	I-21
I.3.1 Inventory for Waterborne Radioactive Materials	I-21
I.3.1.1 Radionuclide Inventory Used in the Model of Long-Term Performance for Proposed Action	I-25
I.3.1.2 Radionuclide Inventory Used in the Model of Long-Term Performance for Inventory Module 1	I-25
I.3.1.3 Radionuclide Inventory Used in the Model of Long-Term Performance for Inventory Module 2	I-26
I.3.2 Inventory for Waterborne Chemically Toxic Materials	I-28
I.3.3 Inventory for Atmospheric Radioactive Materials	I-29
I.4 Extension of TSPA Methods and Models for EIS Analysis of Long-Term Performance	I-29
I.4.1 Methodology	I-30
I.4.2 Assumptions	I-31
I.4.3 Use of Computer Software and Models	I-32
I.4.4 Modifications to the TSPA–Site Recommendation and Supplemental Science and Performance Analysis Models	I-32
I.4.4.1 Modifications to FEHM Particle Tracker Input and Output	I-33

<u>Section</u>	<u>Page</u>
I.4.4.2 Estimation of the Thermal Profiles and Infiltration for the Lower Block	I-34
I.4.4.3 Saturated Zone Breakthrough Curves	I-35
I.4.4.4 Modification to the Waste Package Degradation Model	I-37
I.4.4.5 Early Waste Package Failure	I-38
I.4.4.6 Biosphere Dose Conversion Factors for the 40 CFR 197 Reasonably Maximally Exposed Individual	I-39
I.4.4.7 Igneous Activity Scenario	I-39
I.4.4.8 Human Intrusion Scenario	I-41
I.4.5 Extension of Groundwater Impacts to Other Distances	I-42
I.4.5.1 Locations for Assessing Postclosure Impacts to Human Health	I-42
I.4.5.2 Scaling Approach	I-45
I.4.5.3 Scaling Factors for Dose or Water Quality Concentrations at Longer Distances	I-46
I.5 Waterborne Radioactive Material Impacts	I-46
I.5.1 Waste Package Failure	I-47
I.5.2 Annual Individual Dose for 10,000 Years After Closure	I-48
I.5.3 Annual Individual Dose for 1,000,000 Years After Closure	I-48
I.5.3.1 Annual Individual Dose for the Proposed Action Inventory, Higher- and Lower-Temperature Repository Operating Modes	I-49
I.5.3.2 Annual Individual Dose for Inventory Module 1, Higher-Temperature Repository Operating Mode	I-50
I.5.3.3 Annual Individual Dose for Inventory Module 2, Higher-Temperature Repository Operating Mode	I-50
I.5.3.4 Annual Individual Dose for Igneous Activity Scenario, Higher- and Lower- Temperature Repository Operating Modes	I-50
I.5.3.5 Annual Individual Dose for the Human Intrusion Scenario	I-51
I.5.4 Comparison to Groundwater Protection Standards	I-51
I.6 Waterborne Chemically Toxic Material Impacts	I-52
I.6.1 Screening Analysis	I-52
I.6.1.1 Maximum Source Concentrations of Chemically Toxic Materials in the Repository	I-52
I.6.1.2 Further Screening for Chemically Toxic Materials	I-54
I.6.2 Bounding Consequence Analysis for Chemically Toxic Materials	I-54
I.6.2.1 Assumptions	I-55
I.6.2.2 Surface Area Exposed to General Corrosion	I-56
I.6.2.3 General Corrosion Rates	I-59
I.6.2.4 Release Rates	I-60
I.6.2.5 Summary of Bounding Impacts	I-60
I.7 Atmospheric Radioactive Material Impacts	I-62
I.7.1 Carbon-14 Releases to the Atmosphere	I-63
I.7.2 Atmosphere Consequences to the Local Population	I-64
I.7.3 Screening Argument for Radon	I-66
References	I-85

LIST OF TABLES

<u>Table</u>		<u>Page</u>
I-1	Average net infiltration rates over the unsaturated zone flow and transport model domain for the present-day, monsoon, and glacial transition climate states	I-10
I-2	Average net infiltration rates over the unsaturated zone flow and transport model domain for full-glacial climate states	I-10
I-3	Poisson probabilities for improper heat treatment of waste packages	I-15
I-4	Modeled number of idealized waste packages by category type for the abstracted inventories of the Proposed Action, Inventory Module 1, and Inventory Module 2	I-25
I-5	Abstracted inventory of radionuclides passing the screening analysis in each idealized waste package for commercial spent nuclear fuel, DOE spent nuclear fuel, and high-level radioactive waste for Proposed Action, Inventory Module 1, and Inventory Module 2	I-26
I-6	Greater-Than-Class-C low-level waste volumes by source	I-27
I-7	Abstracted inventory of radionuclides passing the screening analysis in each idealized waste package for Greater-Than-Class-C and Special-Performance-Assessment-Required waste under Inventory Module 2	I-28
I-8	Inventory of chemical materials placed in the Proposed Action repository	I-29
I-9	Carbon-14 gaseous inventory from commercial spent nuclear fuel	I-30
I-10	Average values of the dietary and lifestyle attributes for the RMEI	I-39
I-11	Biosphere dose conversion factors for the RMEI for the groundwater release and the volcanic release exposure scenarios	I-40
I-12	Groundwater impact distance scale factors for 99-percent captured mass, longitudinal dispersivity 100 meters, horizontal dispersivity 10 meters, and vertical dispersivity 0.1 meters	I-47
I-13	Peak mean annual individual doses for analyzed inventories, scenarios, and temperature operating modes	I-48
I-14	Peak 95th-percentile annual individual doses for analyzed inventories, scenarios, and temperature operating modes	I-49
I-15	Peak mean gross alpha activity for analyzed inventories, scenarios, and temperature operating modes	I-53
I-16	Peak mean total radium concentration for analyzed inventories, scenarios, and temperature operating modes	I-53
I-17	Peak mean annual whole body or organ dose for the sum of all beta- and photon-emitting radionuclides during 10,000 years after closure for analyzed inventories, scenarios, and temperature operating modes	I-53
I-18	Total exposed surface area of the Alloy-22 outer layer of all waste packages under the Proposed Action inventory	I-56
I-19	Total exposed surface area of the Alloy-22 rails for all drip shields under the Proposed Action inventory	I-56
I-20	Total exposed surface area of the Alloy-22 outer layer of all waste packages for the Module 1 inventory	I-57
I-21	Total exposed surface area of the Alloy-22 rails for all drip shields for the Module 1 inventory	I-57
I-22	Total exposed surface area of the Alloy-22 outer layer of all waste packages for the Module 2 inventory	I-57
I-23	Total exposed surface area of the Alloy-22 rails for all drip shields for the Module 2 inventory	I-58

<u>Table</u>		<u>Page</u>
I-24	Total exposed surface area of the Alloy-22 components for all emplacement pallets under the Proposed Action, Module 1, and Module 2 inventories	I-58
I-25	Total exposed surface area of the stainless-steel 316NG components for all emplacement pallets under the Proposed Action inventory	I-59
I-26	Total exposed surface area of the stainless-steel 316NG components for all emplacement pallets for the Module 1 inventory	I-59
I-27	Total exposed surface area of the stainless-steel 316NG components for all emplacement pallets for the Module 2 inventory	I-59
I-28	Bounding mass release rates from Alloy-22 and stainless-steel 316NG components from general corrosion for the Proposed Action	I-60
I-29	Bounding mass release rates from Alloy-22 and stainless-steel 316NG components from general corrosion for Module 1	I-60
I-30	Bounding mass release rates from Alloy-22 and stainless-steel 316NG components from general corrosion for Module 2	I-61
I-31	Bounding concentrations of waterborne chemical materials of concern compared to Maximum Contaminant Levels Goals	I-61
I-32	Summary of intake of waterborne chemical materials of concern based on maximum bounding concentrations listed in Table I-31 compared to Oral Reference Doses	I-62
I-33	Meteorologic joint frequency data used for Yucca Mountain atmospheric releases	I-65

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
I-1	TSPA model	I-68
I-2	Development of abstracted inventory for TSPA-FEIS	I-69
I-3	Approximate configuration of the potential Yucca Mountain Repository	I-70
I-4	The four saturated zone capture regions in relation to the primary repository block for the Proposed Action	I-71
I-5	The four saturated zone capture regions in relation to the primary and lower repository blocks for Inventory Modules 1 and 2	I-72
I-6	Outline of the Lower Block showing the locations of the 51 particle-tracking nodes	I-73
I-7	Southernmost boundary of the controlled area and the accessible environment	I-74
I-8	Waste package failure curves for 300 probabilistic simulations for the Proposed Action inventory; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of these simulations	I-75
I-9	Cladding failure profile for the Proposed Action inventory	I-75
I-10	Comparison plot of the total mean annual individual dose at the RMEI location under the higher-temperature and lower-temperature operating modes for the Proposed Action inventory, nominal scenario	I-76
I-11	Total and individual radionuclide mean annual dose to an individual at the RMEI location for the higher-temperature operating mode for the Proposed Action inventory, nominal scenario	I-76
I-12	Total annual individual dose at the RMEI location for 300 probabilistic simulations of the higher-temperature operating mode for the Proposed Action inventory, nominal scenario; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of these simulations	I-77

<u>Figure</u>		<u>Page</u>
I-13	Total annual individual dose at the RMEI location for 300 probabilistic simulations of the lower-temperature operating mode for the Proposed Action inventory, nominal scenario; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of these simulations	I-77
I-14	Total annual individual dose at the RMEI location for 300 probabilistic simulations of the higher-temperature operating mode for the Module 1 inventory, nominal scenario; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of these simulations	I-78
I-15	Total annual individual dose at the RMEI location for 300 probabilistic simulations of the higher-temperature operating mode for the Module 2 incremental inventory, nominal scenario; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of these simulations	I-78
I-16	Comparison plot of the mean total annual individual dose at the RMEI location for the higher-temperature operating mode for the Proposed Action, Module 1, and incremental Module 2 inventories, nominal scenario	I-79
I-17	Total annual individual dose at the RMEI location for 500 out of 5,000 probabilistic simulations of the higher-temperature operating mode for the Proposed Action inventory under the igneous activity scenario; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of all 5,000 simulations	I-79
I-18	Total mean individual receptor dose at the RMEI location for the higher-temperature operating mode for the Proposed Action inventory under the igneous activity scenario; the figure displays the mean results for both the eruptive and intrusive events and the sum of these events as “Total Igneous”	I-80
I-19	Total mean annual individual dose at the RMEI location for the higher-temperature and lower-temperature operating modes for the Proposed Action inventory under the igneous activity scenario	I-80
I-20	Total annual individual dose at the RMEI location for 500 out of 5,000 probabilistic simulations of the lower-temperature operating mode for the Proposed Action inventory under the igneous activity scenario; the figure also displays the 5th-percentile, median, mean, and 95th-percentile values of these simulations	I-81
I-21	Total mean annual individual dose at the RMEI location for the lower-temperature operating mode for the Proposed Action inventory under the igneous activity scenario; the figure displays the mean results for both the eruptive and intrusive events and the sum of these events as “Total Igneous”	I-81
I-22	Total annual individual dose at the RMEI location for 300 probabilistic simulations of the higher-temperature operating mode for the Proposed Action inventory under the human-intrusion-at-30,000-years scenario; the figure also displays the median, mean, and 95th-percentile values of these simulations	I-82
I-23	Mean activity concentrations of gross alpha activity and total radium (radium-226 plus radium-228) at the RMEI location of 300 probabilistic simulations of the higher-temperature operating mode for the Proposed Action inventory for the nominal scenario	I-82
I-24	Mean activity concentrations of gross alpha activity and total radium (radium-226 plus radium-228) at the RMEI location of 300 probabilistic simulations of the lower-temperature operating mode for the Proposed Action inventory for the nominal scenario	I-83

<u>Figure</u>		<u>Page</u>
I-25	Mean dose to critical organs for technetium-99, carbon-14, and iodine-129 at the RMEI location of 300 probabilistic simulations of the higher-temperature operating mode for the Proposed Action inventory for the nominal scenario	I-83
I-26	Mean dose to critical organs for technetium-99, carbon-14, and iodine-129 at the RMEI location of 300 probabilistic simulations of the lower-temperature operating mode for the Proposed Action inventory for the nominal scenario	I-84
I-27	Fraction of perforated cladding for commercial spent nuclear fuel as a function of time after repository closure	I-84
I-28	Release rate of carbon-14 from the repository to the ground surface for 80,000 years following repository closure	I-85